

A Novel Unstable Duplication Upstream of *HAS2* Predisposes to a Breed-Defining Skin Phenotype and a Periodic Fever Syndrome in Chinese Shar-Pei Dogs

Mia Olsson^{1*}, Jennifer R. S. Meadows^{1,9}, Katarina Truvé^{2,9}, Gerli Rosengren Pielberg^{1,9}, Francesca Puppo^{3,9}, Evan Mauceli⁴, Javier Quilez⁵, Noriko Tonomura⁴, Giordana Zanna⁶, Maria José Docampo⁷, Anna Bassols⁷, Anne C. Avery⁸, Elinor K. Karlsson^{4,9}, Anne Thomas¹⁰, Daniel L. Kastner³, Erik Bongcam-Rudloff¹¹, Matthew T. Webster¹, Armand Sanchez⁵, Åke Hedhammar¹², Elaine F. Remmers³, Leif Andersson^{1,2}, Lluís Ferrer⁶, Linda Tintle^{13*}, Kerstin Lindblad-Toh^{1,4*}

1 Science for Life Laboratory, Department of Medical Biochemistry and Microbiology, Uppsala University, Uppsala, Sweden, **2** Department of Animal Breeding and Genetics, Swedish University of Agricultural Sciences, Uppsala, Sweden, **3** National Human Genome Research Institute, Bethesda, Maryland, United States of America, **4** Broad Institute of Harvard and Massachusetts Institute of Technology (MIT), Cambridge, Massachusetts, United States of America, **5** Department of Animal and Food Science, Veterinary Molecular Genetics Service, Universitat Autònoma de Barcelona, Barcelona, Spain, **6** Department of Animal Medicine and Surgery, Universitat Autònoma de Barcelona, Barcelona, Spain, **7** Department of Biochemistry and Molecular Biology, Universitat Autònoma de Barcelona, Barcelona, Spain, **8** Department of Microbiology, Immunology, and Pathology, Colorado State University, Fort Collins, Colorado, United States of America, **9** FAS Center for Systems Biology, Harvard University, Cambridge, Massachusetts, United States of America, **10** ANTAGENE Laboratory, Limonest, France, **11** Linnaeus Centre for Bioinformatics, Uppsala University, Uppsala, Sweden, **12** Department of Clinical Sciences, Swedish University of Agricultural Sciences, Uppsala, Sweden, **13** Wurtsboro Veterinary Clinic, Wurtsboro, New York, United States of America

Abstract

Hereditary periodic fever syndromes are characterized by recurrent episodes of fever and inflammation with no known pathogenic or autoimmune cause. In humans, several genes have been implicated in this group of diseases, but the majority of cases remain unexplained. A similar periodic fever syndrome is relatively frequent in the Chinese Shar-Pei breed of dogs. In the western world, Shar-Pei have been strongly selected for a distinctive thick and heavily folded skin. In this study, a mutation affecting both these traits was identified. Using genome-wide SNP analysis of Shar-Pei and other breeds, the strongest signal of a breed-specific selective sweep was located on chromosome 13. The same region also harbored the strongest genome-wide association (GWA) signal for susceptibility to the periodic fever syndrome ($p_{\text{raw}} = 2.3 \times 10^{-6}$, $p_{\text{genome}} = 0.01$). Dense targeted resequencing revealed two partially overlapping duplications, 14.3 Kb and 16.1 Kb in size, unique to Shar-Pei and upstream of the *Hyaluronic Acid Synthase 2* (*HAS2*) gene. *HAS2* encodes the rate-limiting enzyme synthesizing hyaluronan (HA), a major component of the skin. HA is up-regulated and accumulates in the thickened skin of Shar-Pei. A high copy number of the 16.1 Kb duplication was associated with an increased expression of *HAS2* as well as the periodic fever syndrome ($p < 0.0001$). When fragmented, HA can act as a trigger of the innate immune system and stimulate sterile fever and inflammation. The strong selection for the skin phenotype therefore appears to enrich for a pleiotropic mutation predisposing these dogs to a periodic fever syndrome. The identification of HA as a major risk factor for this canine disease raises the potential of this glycosaminoglycan as a risk factor for human periodic fevers and as an important driver of chronic inflammation.

Citation: Olsson M, Meadows JRS, Truvé K, Rosengren Pielberg G, Puppo F, et al. (2011) A Novel Unstable Duplication Upstream of *HAS2* Predisposes to a Breed-Defining Skin Phenotype and a Periodic Fever Syndrome in Chinese Shar-Pei Dogs. *PLoS Genet* 7(3): e1001332. doi:10.1371/journal.pgen.1001332

Editor: Michel Georges, University of Liège, Belgium

Received: June 3, 2010; **Accepted:** February 10, 2011; **Published:** March 17, 2011

This is an open-access article distributed under the terms of the Creative Commons Public Domain declaration which stipulates that, once placed in the public domain, this work may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose.

Funding: This work was supported by the Swedish Research Council; FORMAS; the Swedish Research Council for Environment, Agricultural Sciences, and Spatial Planning; the Swedish Foundation for Strategic Research; in part by the Intramural Research Program of the National Institute of Arthritis and Musculoskeletal and Skin Diseases and the National Human Genome Research Institute of the National Institutes of Health, the Chinese Shar-Pei Charitable Trust and the European Commission (FP7-LUPA, GA-201370). KL-T is the recipient of a EURYL award from the European Science Foundation. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors KL-T, MO, and LT have filed a patent for development of a genetic test.

* E-mail: mia.olsson@imbim.uu.se (MO); wvc@warwick.net (LT); kersli@broadinstitute.org (KL-T)

† These authors contributed equally to this work.

Introduction

Shar-Pei dogs have been companion animals for centuries within China where they were commissioned to guard and hunt, and to sometimes serve as fighting animals. At the beginning of the communist era dog ownership was highly taxed and the breed was

brought close to extinction. A few Chinese Shar-Pei dogs were exported to the United States in the early 1970's and Shar-Pei descending from this limited number of animals have undergone strong selection for a wrinkled skin phenotype and heavily padded muzzle and are called the "meatmouth" type (Figure 1A–1C) and have now found global popularity. The ancestral Shar-Pei,

Author Summary

Shar-Pei dogs have two unique features: a breed defining “wrinkled” skin phenotype and a genetic disorder called Familial Shar-Pei Fever (FSF). The wrinkled phenotype is strongly selected for and is the result of excessive hyaluronan (HA) deposited in the skin. HA is a molecule that may behave in a pro-inflammatory manner and create a “danger signal” by being analogous to molecules on the surface of pathogens. FSF is characterized by unprovoked episodes of fever and/or inflammation and resembles several human autoinflammatory syndromes. Here we show that the two features are connected and have the same genetic origin, a regulatory mutation located close to a HA synthesizing gene (*HAS2*). The mutation is a 16.1 Kb duplication, the copy number of which correlates with *HAS2* expression and disease. We suggest that the large amount of HA responsible for the skin condition predisposes to sterile fever and inflammation. *HAS2* was previously not known to associate with autoinflammatory disease, and this finding is of wide interest since approximately 60% of human patients with periodic fever syndrome remain genetically unexplained. This investigation also demonstrates how strong artificial selection may affect not only desired and selected phenotypes, but also the health of domestic animals.

referred to as the “traditional” type Shar-Pei, still occurs and it presents with a less accentuated skin condition (Figure 1D). The major constituent of the deposit in the thickened skin is hyaluronan or hyaluronic acid (HA). HA is a large, multifunctional, linear, negatively charged, non-sulfated glycosaminoglycan of the extracellular and pericellular matrices. It is composed of repeating disaccharides and is widely spread throughout epithelial, connective and neural tissues [1,2]. The biological role of HA depends on its size, location and equilibrium between synthesis and degradation [1–3]. Meatmouth Shar-Pei show two- to five-fold higher serum levels of HA compared to other breeds [4], allowing us to propose the term *hyaluronanosis*, a definition also used for a comparable human condition [5]. HA is synthesized at the plasma membrane by three HA synthases, HAS1, HAS2 and HAS3, with HAS2 being the rate limiting-enzyme [6]. *HAS2* is

overexpressed in dermal fibroblasts of Shar-Pei compared with other canine breeds [7] suggesting a regulatory mutation as causative for hyaluronanosis. HA is deposited throughout the skin of Shar-Pei, often in microscopic lakes and grossly evident vesicles, leading to the formation of thickened skin folds around the head and tibiotarsal (hock) joints (Figure 1E). Almost all Shar-Pei seem to be affected by hyaluronanosis, however the extent varies among individuals and adults exhibit less skin folds and hyaluronanosis than puppies. Strong selection by breeders for dogs who retained their skin folds into adulthood has altered the phenotype of the breed to the more commonly heavily wrinkled meatmouth type.

Meatmouth Shar-Pei also suffer a strong predisposition to an autoinflammatory disease, Familial Shar-Pei Fever (FSF), which clinically resembles some human hereditary periodic fever syndromes, such as Familial Mediterranean Fever (FMF) [8]. Both diseases are characterized by seemingly unprovoked episodes of fever and inflammation and both FMF and FSF present as short (12–48 hour) recurrent bouts of high fever, accompanied by localized inflammation usually involving major joints (especially the tibiotarsal joints). Patients with FMF or Shar-Pei with FSF can suffer episodes as often as every few weeks, but in the interim seem symptom free. However, since acute phase reactants may endure between episodes, a subclinical state and chronic autoinflammation may persist (Linda Tintle unpublished data). As a secondary complication, the chronic state puts human patients, as well as affected Shar-Pei dogs, at risk of developing reactive systemic AA amyloidosis and subsequent kidney or liver failure [8,9]. In Shar-Pei, the fever episodes are typically more frequent during the first years of life and the percentage of affected dogs is very high, estimated to be 23% in the US in 1992 [9].

Results

In order to find candidate loci for the breed-specific phenotype (hyaluronanosis), known to be under selective pressure, we screened the genome for signatures of selective sweeps. These sweeps can be recognized as long chromosomal segments with a low degree of heterozygosity within populations [10]. Using 50,000 single nucleotide polymorphisms (SNPs) distributed throughout the dog genome, the level of heterozygosity in windows of ten consecutive SNPs was compared between a set




Figure 1. The phenotypic spectrum of the Chinese Shar-Pei. Following strong selection for the “wrinkled” skin phenotype, Shar-Pei dogs in the western world most commonly present as the meatmouth type (A–C). The traditional type of Shar-Pei (D) is the ancestral version and is still common in China. The characteristic skin is a result of a deposition of mucin, mainly hyaluronic acid (HA), in the upper dermis of the skin. The deposit collects in certain areas of Shar-Pei skin and often as “socks” around the hocks (E). The meatmouth Shar-Pei (A–C) is also predisposed to a breed-specific periodic fever syndrome called Familial Shar-Pei Fever (FSF).
doi:10.1371/journal.pgen.1001332.g001

of Shar-Pei ($n = 50$, all from the US, Table S1) and the average of 24 other canine breeds ($n = 230$). On four chromosomes (*Cfa* 5, 6, 13 and X) the reduction in heterozygosity in Shar-Pei was greater than 4-fold the average of control breeds (Figure 2A). The strongest signal of reduced heterozygosity appeared within a 3.7 Mb stretch on chromosome 13 (CanFam 2.0 Chr13: 23,487,992–27,227,623) (<http://genome.ucsc.edu/>) near the *HAS2* gene, where almost complete homozygosity was observed in Shar-Pei (Figure 2C). Here the reduction in heterozygosity was greater than 10-fold in Shar-Pei and several smaller regions showed complete homozygosity. The same region was confirmed to show high levels of homozygosity when the analysis was repeated in 37 additional Shar-Pei dogs sampled from Spain (Table S1) and was overlapping a sweep region reported by others for this breed [11]. The strong signal, together with the known function of *HAS2* and its aberrant expression pattern in Shar-Pei, made this region an obvious candidate for the mutation causing the wrinkled skin phenotype (hyaluronanosis).

In parallel, we performed a genome-wide association study to map the susceptibility locus for FSF, using Shar-Pei strictly classified as FSF affected ($n = 24$, classification code FSF+A and FSF+, described in Materials and Methods) and unaffected ($n = 17$, classification code H+, described in Materials and Methods). Five SNPs were significantly associated (best SNP $p_{\text{raw}} = 7.0 \times 10^{-7}$, $p_{\text{genome}} = 0.005$ based on 100,000 permutations; software package PLINK <http://pngu.mgh.harvard.edu/~purcell/plink> [12]), all on chromosome 13 (CanFam 2.0 Chr13: 22.4–30.7 Mb, Figure 2B). After correcting for putative stratification, two outlier cases were removed (Figure S1) and the same SNPs, forming the same signal of association remained (best SNP $p_{\text{raw}} = 2.3 \times 10^{-6}$, $p_{\text{genome}} = 0.01$; Table S2) with a genomic inflation factor of 1.2. When the association signal and the sweep signal were compared they appeared interspersed, so that individual SNPs were either part of homozygous regions or showed association with FSF (Figure 2C). It was therefore difficult to determine exactly where the strongest association fell, as variation is required to detect association.

Targeted sequence capture technology was used to further investigate the sweep signal and to search for the hyaluronanosis causative mutation. We resequenced 1.5 Mb around and upstream of our candidate gene, *HAS2* (CanFam 2.0 Chr13: 22,937,592–24,414,650) in four Shar-Pei (two meatmouth type with high serum HA levels and two traditional type) and three control dogs from other breeds. The obtained sequences were mapped to the boxer reference sequence providing at least 5X coverage for 96–98% of the resequenced region in each individual. The targeted region also included the large intergenic noncoding RNA, *HAS2 antisense (HAS2as)*; Table S3) which has been proposed as a negative post-transcriptional regulator of *HAS2* mRNA [13]. After masking repetitive sequences we identified ~670 indels and ~1,500 SNP in each dog (Table S4) as well as two overlapping duplications in the Shar-Pei (Figure 3A). Nine mutations (eight SNPs and one indel) located in conserved elements as well as two SNPs possibly regulating transcription, were selected for further investigation due to their unique pattern in the sequenced Shar-Pei dogs. Additional genotyping in Shar-Pei and dogs from other breeds (Tables S1, S5) showed these mutations were not specific to Shar-Pei and the variants were subsequently excluded as causative.

The two duplications were named after the Shar-Pei type in which they were first identified. The “meatmouth” duplication was the larger fragment, 16.1 Kb (CanFam 2.0 Chr13: 23,746,089–23,762,189) with breakpoints located in repeats (a SINE at the centromeric end and a LINE at the telomeric end) and individual copies separated by seven base pairs (Figure 3B). The “traditional” duplication was 14.3 Kb (CanFam 2.0 Chr13:

23,743,906–23,758,214) and was identified in the two Shar-Pei with a less accentuated skin phenotype (Figure 3B). We first examined the duplications via Southern blot with control breeds ($n = 2$), traditional ($n = 2$) and meatmouth Shar-Pei ($n = 6$) (Figure 3C). As the digest cut outside and within both duplications, we were able to observe the absence of the variants from control breeds and separate restriction patterns in traditional and meatmouth type Shar-Pei. Interestingly, one meatmouth dog contained both duplication types (Figure 3C lane 6 and confirmed by PCR across break points, data not shown). Two copy number assays were developed to quantify these elements. The first (CNV-E) measured only the meatmouth duplication whilst the second (CNV-748), detected both the traditional and meatmouth duplications. Copy number analysis was estimated as the relative fold enrichment ($\Delta\Delta\text{Ct}$) between an amplicon within the duplication and one outside the duplication in a housekeeping gene. Assay CNV-E was run on 90 Shar-Pei and 73 dogs from 24 other breeds (Table S1) and assay CNV-748 on a subset of 44 Shar-Pei and 14 dogs from other breeds. Assay CNV-748 demonstrated that both the traditional and meatmouth duplications are unique to the Shar-Pei breed (Figure 4 and Figure S2).

We used the results of both assays to search for a relationship between Familial Shar-Pei Fever (FSF) and either meatmouth copy number (Assay CNV-E), traditional copy number (the normalized difference between CNV-748 and CNV-E) or total traditional+meatmouth copy number (Assay CNV-748). Shar-Pei dogs were strictly classified as affected by FSF ($n = 28$, FSF+A and FSF+) or unaffected by FSF ($n = 16$, H+). The most significant association was found when only the meatmouth copy number was considered ($p < 0.0001$, Figure 4) although a weaker association with total copy number ($p < 0.01$) was also seen. The observed association between fever and meatmouth copy number, despite the very high homozygosity in this region, strongly suggests that a high copy number is not just a genetic marker for FSF but is causally related to the development of disease.

Of the 153 dogs analyzed with the meatmouth copy number assay, 31 Shar-Pei and 18 control animals also had serum measures of HA available. No clear association was detected between HA levels and copy number (Figure S3), however the mean HA level in Shar-Pei with \geq six copies was 905 ± 403 ug/L ($n = 21$), whilst Shar-Pei with fewer copies had a mean concentration of 770 ± 494 ug/L ($n = 12$) and control breeds had HA serum levels of 206 ± 145 ug/L ($n = 19$). Interestingly, the three traditional Shar-Pei dogs had serum HA levels between 73 and 266 ug/L, which fell within the normal range [4].

The link between copy number and the expression of *HAS2* and *HAS2as* was examined on a smaller scale using dermal fibroblasts cultured from six separate meatmouth Shar-Pei. The expression of both genes was calibrated against the Shar-Pei with lowest copy number (CNV estimate = 5) and both genes showed an increasing trend of expression with copy number (Figure 5). These data suggest that a regulatory element for *HAS2* is located in the duplicated region, however the interpretation of the *HAS2as* result is less clear. A single study of a human osteosarcoma cell line demonstrated that the expression of two isoforms of *HAS2as* were able to reduce *HAS2* expression, and so these mRNAs may act as regulators of HA production [13]. Our data could indicate that *HAS2as* expression is also influenced by a regulator element in the duplication, or that *HAS2as* is up-regulated in response to *HAS2* levels. If either of these scenarios were true, it is possible that if RNA expression were measured at multiple time points we would see temporal *HAS2* repression. It could also be that the interaction between canine fibroblast *HAS2* and *HAS2as* does not mirror the human system and that the canine antisense mRNA is non-




Figure 2. The association with Shar-Pei Fever susceptibility and the strongest selective sweep signal co-localize on chromosome 13. (A) A 10-fold reduction of heterozygosity was identified on chromosome 13 when comparing Shar-Pei ($n = 50$) to 24 other canine breeds ($n = 230$). 50,000 SNPs were used to screen the whole genome using a sliding window approach (see Materials and Methods). (B) A case-control genome-wide

association analysis identified a strong peak with several SNPs on chromosome 13 to be in association with Familial Shar-Pei Fever (FSF). After correcting for stratification and multiple testing (100,000 permutations), four SNPs retained significant association ($p < 0.05$; strongest SNP association, CanFam 2.0 chr13: 27,913,803 Mb). Shar-Pei dogs used in the study were strictly classified into groups of affected ($n = 22$) and unaffected ($n = 17$) by FSF. (C) SNPs associated with FSF (blue line) are interspersed with the signals of selection (red line). The 39 Shar-Pei and 17,227 SNP common to both analyses were used to generate this graph.
doi:10.1371/journal.pgen.1001332.g002

functional. At present our results must be considered as preliminary and it is clear that further exploration of the interaction between canine *HAS2* and *HAS2as* is required.

Discussion

Here we have identified a 16.1 Kb duplication located approximately 350 Kb upstream of *HAS2*. This is clearly a derived mutation since it occurs as a single copy sequence in other dog breeds. We postulate that this is a causative mutation associated with both hyaluronanosis and Shar-Pei fever, as the observed correlation between copy number and susceptibility to Shar-Pei fever was not expected if this was a linked, neutral polymorphism. We suggest that the unique region of the meatmouth type duplication identified in Shar-Pei contains one or more regulatory elements that alter the expression of *HAS2*. It appears possible that as the duplication copy number increases, so does the copy number of potential enhancer elements within the duplication, likely leading to a higher expression of *HAS2* and elevated HA levels, and resulting in the development of hyaluronanosis in this breed. We propose a scenario whereby the traditional duplication arose *de novo* in the traditional type of Shar-Pei causing a milder skin phenotype. This event made the region unstable and allowed the second meatmouth duplication to occur. Breeders subsequently selected the meatmouth duplication as a higher copy number enhanced the phenotypic effect in appearance. However, it is not yet possible to say whether the meatmouth duplication first occurred at low frequency in the Chinese Shar-Pei population and quickly rose during breeding in America, or if the mutation occurred spontaneously during breed expansion in the West.

Tandem duplications are notoriously unstable and may show copy number variation due to unequal crossing-over, as is clearly illustrated by the copy number variation of a 450 Kb duplication associated with dominant white colour in pigs [14]. The meatmouth Shar-Pei duplication adds to the list of copy number variants (CNVs), which affect phenotypic traits in domestic animals (e.g. dominant white in pigs [14], gray color in horses [15], the hair-ridge in Rhodesian ridgeback dogs [16], and pea-comb in chicken [17]), several of which are linked not only to the desirable trait but also to disease. Interestingly, all of these except pea-comb, represent novel duplications derived from single copy sequences. This is in contrast to most reported CNVs in humans, which are mainly benign and represent expansions or contractions of duplicated sequences [18].

Although we failed to find a significant correlation between serum HA levels and copy number, this does not exclude our proposed hyaluronanosis scenario. Difficulties in correlating fluctuating serum levels of HA with other clinical and biomedical parameters have also been reported in many human studies, where no or only weak correlations were observed [19,20]. We have shown that the 16.1 Kb duplication appears only in meatmouth Shar-Pei, a breed type that has elevated levels of HA compared to both traditional Shar-Pei and other breeds, and that copy number correlates with a breed-specific syndrome associated with excessive HA deposition and the over expression of a HA synthesizing gene. Because HA is primarily a component of

the extracellular matrix, serum measurements may only broadly reflect total body HA.

Hyaluronan can bind to several cellular receptors (e.g. CD44, RHAMM and layilin), however it is the interaction between CD44 and HA which acts as a biological regulator, differentially modulating the cellular microenvironment in response to homeostatic versus inflammatory conditions [21]. Alterations in the balance between native high molecular weight HA versus fragmented HA may result in activation of innate immunity. HA has been linked to sterile inflammation as an endogenous response molecule to sterile tissue injury [21]. Shorter fragments of HA can be generated by environmental insults such as sterile trauma [22], reactive oxidative species (ROS) [23], or pathogenic hyaluronidases, and it is these low molecular weight fractions which can become pro-inflammatory danger associated molecular pattern (DAMP) molecules [22,24] mimicking microbial surface molecules.

Using a mouse model, Yamasaki and colleagues [25] showed that HA can interact with the cell through two separate pathways that culminate in the release of IL-1 β , which together with IL-6, is one of the main promoters of fever. In the first route, CD44 bound HA is degraded at the plasma membrane by hyaluronidase-2 (HYAL2) prior to endocytosis and further cleavage by lysosomal hyaluronidase-1 (HYAL1). The resultant small intracellular oligosaccharides of HA activate the NLRP3 inflammasome, a multiprotein complex consisting of the NLRP3 scaffold, the ASC adaptor and caspase-1 [26]. In the second arm, the CD44-HA complex activates toll like receptors 2 and 4 (TLR2 and 4), leading to intracellular IL-1 β mRNA transcription and the formation of pro-IL-1 β . Activation of the NLRP3 inflammasome by HA oligosaccharides allows cleavage of this pro-IL-1 β by caspase-1 and subsequent release of IL-1 β . The NLRP3 inflammasome is present in the cytosol of many cells including monocytes, macrophages and mast cells, and has been implicated in the pathogenesis of numerous autoinflammatory diseases in humans including the cryopyrin-associated periodic syndromes which result from mutations in *NLRP3/CIAS1* [26].

The actual role of excessive HA in Shar-Pei needs to be investigated further. Shar-Pei may experience exogenous fragmentation of their over-abundant HA from sterile or pathogenic trauma. This, plus endogenous degradation of excessive native HA, may contribute to induction of recurrent episodes of fever and inflammation. Acute fever events in Shar-Pei respond rapidly to dipyrone, a potent antipyretic and analgesic pyrazolone, which has been demonstrated to inhibit IL-1 β induced fever [27–29 and Linda Tintle unpublished data]. It is therefore not surprising that the strong selection on the hyaluronanosis phenotype, with increased levels of cutaneous HA, may predispose Shar-Pei to autoinflammation, potentially contributing to other pathologies seen in this breed. One such example is renal medullary amyloidosis. Histopathologically, kidneys of Shar-Pei in renal failure have multifocal non-suppurative tubulointerstitial nephritis with fibrosis. Medullary amyloidosis predominates and glomerular deposition, although consistent, is highly variable in its extent [8,30]. The renal medulla is naturally HA rich and enhanced renal interstitial HA accumulation can be coupled to inflammatory responses, such as ischemia-reperfusion injury, transplant-rejec-




Figure 3. The identification of two breed-specific duplications in Shar-Pei. (A) Targeted resequencing of a 1.5 Mb region on chromosome 13 identified a duplication with on average 3.5–4.5X higher read coverage in two meatmouth Shar-Pei (black and red), compared to three control breeds (green, Standard Poodle; orange, Neapolitan Mastiff and purple, Pug). A shorter duplication was detected in the traditional Shar-Pei (blue). (B) The meatmouth duplication was determined to be 16.1 Kb long (CanFam 2.0 Chr13: 23,746,089–23,762,189) with both breakpoints located in repeats (a SINE and a LINE) and with an insertion of 7 bp separating different copies. The duplication in the traditional Shar-Pei overlapped the meatmouth duplication and was slightly shorter, 14.3 Kb long (CanFam 2.0 Chr13: 23,743,906–23,758,214). In this case the copies were separated by 1 bp but were still anchored in repeat motifs (c) Southern blot analysis using *Bsr*GI digested gDNA from Shar-Pei and control breeds confirmed the existence of two duplication types in Shar-Pei. One meatmouth dog (lane 6) contained both duplication types. Individuals were classified as healthy (h) or as affected by Familial Shar-Pei Fever (f). doi:10.1371/journal.pgen.1001332.g003

tion, tubulointerstitial inflammation and diabetes [31]. In addition, Shar-Pei are prone to mast cell disease including mast cell tumors [32,33]. The binding of HA to CD44 has been shown to play a critical role in regulation of murine cutaneous and connective

tissue mast cell proliferation [34]. As the CD44-HA interaction may modulate local immune responses through regulation of mast cell functions [35], excessive HA and its subsequent damage and degradation may play a role also in the Shar-Pei breed's




Figure 4. The relationship between copy number estimate and susceptibility to Familial Shar-Pei Fever. A significant correlation ($p = < 0.0001$, Mann Whitney test) was seen when the meatmouth copy number in unaffected Shar-Pei ($n = 16$, H+) and individuals affected by FSF ($n = 28$, FSF+ and FSF+ A) were compared. Based on this limited sample size, most dogs with more than six copies had fever whereas most dogs with less than four copies did not.

doi:10.1371/journal.pgen.1001332.g004

predilection for allergic skin disease and other mast cell driven inflammation.

This study suggests that *HAS2* dysregulation can trigger a periodic fever syndrome in dogs and therefore it will be relevant to examine the approximately 60% of human fever patients who currently have unexplained disease. Previously, the role of hyaluronan in sterile inflammation has focused on HA signaling and degradation; for example a deficiency of hyaluronidase causing mucopolysaccharidosis type IX in humans has some autoinflammatory features [36]. However by directly implicating *HAS2* in inflammation, we suggest that a reexamination of genes further up the biosynthetic pathway, such as those involved in HA synthesis and polymerization is called for. In addition, the canine mutation appears regulatory in nature and therefore regulators of HA should be also be included in a broader scope pathway analysis of human patients with unexplained autoinflammatory disease.

Finally, this study illustrates how copy number variations can shape phenotypic traits and how strong artificial selection for certain phenotypic traits may not only affect the desired trait but also the health of the animal.

Materials and Methods

Samples and diagnostic procedure

All dog samples were collected from pet dogs after owner consent following the ethical approval protocols (SLU, Dnr: C103/10, MIT

0910-074-13). DNA was extracted from blood samples using QIAamp DNA Blood Midi Kit (QIAGEN) or PureLink Genomic DNA kit (Invitrogen). All dogs, their breed type, geographic origin, health status and experiment in which they were utilized are listed in Table S1.

Classification of Shar-Pei fever: Purebred Shar-Pei individuals were divided into the following six groups based on their medical records and evidence by owner and/or veterinarian:

1. *FSF+A*, the individual had experienced recurrent episodes of high fever accompanied by inflammation of joints from an early age (less than one year old). Additionally, post-mortem examination detected depositions of amyloid in kidneys and/or liver (amyloidosis).

2. *FSF+*, the individual had experienced recurrent episodes of high fever accompanied with inflammation of joints from an early age (less than one year old).

3. *Atypical FSF*, the individual had experienced occasional unexplained fever episodes or recurrent episodes with a late onset (greater than three years old).

4. *H+*, the individual had never experienced unexplained fever and/or inflammation, was older than five years old at the time of sampling and also lacked first-degree relatives that could be classified into the groups *FSF+A*, *FSF+* or *Atypical FSF*.

5. *H-*, the individual had never experienced unexplained fever and/or inflammation but was younger than 5 years at the time of sampling and/or had first-degree relatives that could be classified into the groups *FSF+A*, *FSF+* or *Atypical FSF*.




Figure 5. Expression analysis reveals a trend of increased *HAS2* and *HAS2as* expression with copy number. Expression levels were measured in dermal fibroblasts that were cultured from individual Shar-Pei skin biopsies. The individual with the lowest copy number (CNV = 5) was used to calibrate each assay.
doi:10.1371/journal.pgen.1001332.g005

6. *Unknown*, the individual's medical record was not available.

Hyaluronanosis: Serum Hyaluronic Acid (HA) concentration was used as a proxy for hyaluronanosis but no distinct cut-off value was established. However, dogs with normal and abnormal concentrations of serum HA were interpreted as before [4]. HA measurements were performed using the Hyaluronan ELISA kit (Echelon Biosciences INC) according to the manufacturer's instructions. The absorbance was read at 405 nm, and a semi-log standard curve was used to calculate hyaluronic acid concentrations.

Homozygosity and genome-wide association mapping

A whole genome scan was performed with two array types, the 27K (v1) and 50K (v2) canine Affymetrix SNP chips. Results were called using Affymetrix's snp5-geno-qc software. The 50K array was used when the rate of heterozygosity was calculated for US Shar-Pei separately and for a reference group of 24 other breeds. The ratio of heterozygosity in 10 SNP (≈ 1 Mb) sliding windows between the two groups was used as a measure of relative heterozygosity. To look for regions of homozygosity within the Shar-Pei genome only, the software package PLINK [12] was used. This was performed both for the 50 K array with 50 US Shar-Pei and replicated for 37 Spanish Shar-Pei using 22,362 SNPs genotyped with the Illumina CanineSNP20 BeadChip. These data were collected with an Illumina BeadStation scanner and genotypes were scored using GenomeStudio. Regions of homozygosity were defined if shared across all Shar-Pei samples.

A case-control association analysis using 17,227 SNP common to both the 27K and 50K arrays (MAF > 0.05, call rate > 75%) was performed in Shar-Pei classified as affected (FSF+A and FSF+, $n = 39$) or unaffected (H+, $n = 17$) by Shar-Pei fever. The software package PLINK [12] was used for the analyses and to ensure

genome-wide significance, p-values were corrected for multiple testing. Values used are the max (T) empirical p-values obtained after 100,000 permutations. To assess whether signals from the two genome scans overlapped, the 39 Shar-Pei with unambiguous phenotypes were analyzed with the 17,227 SNPs common to both SNP platforms.

Targeted resequencing

Targeted capture of the 1.5 Mb candidate region (CanFam 2.0 Chr13: 22,937,592–24,414,650) was performed using a 385K custom-designed sequence capture array from Roche NimbleGen. Hybridization library preparation was performed as following: Genomic DNA (15–20 μ g) was fragmented using sonication; blunting of DNA fragments using T4 DNA Polymerase, Klenow Fragment and T4 Polynucleotide Kinase; adding A-overhangs using Klenow Fragment exo^- and ligation of adaptors using T4 DNA Ligase with Single-read Genomic Adapter Oligo Mix (Illumina). All enzymes were purchased from Fermentas and used following manufacturers instructions. Purification steps were performed using QIAquick PCR Purification Kit (QIAGEN). Hybridization was performed following the manufacturer's instructions without amplification of the fragment library prior to hybridization. Eluted captured DNA and uncaptured libraries were amplified using Phusion High Fidelity PCR Master Mix (Finnzymes) and the SYBR Green PCR Master Mix (Applied Biosystems) was used to estimate the relative fold-enrichment. Capture libraries with the estimated enrichment-factor of >200 were sequenced using Genome Analyzer (Illumina) and obtained sequences were aligned to CanFam 2.0 [37] and to the targeted region using Maq assembly (<http://maq.sourceforge.net/>) [38]. For each individual, sequence coverage was calibrated by dividing the coverage in 100 bp windows by the average coverage for the

total region. Three control breeds (Pug, Neapolitan Mastiff, Standard Poodle) and two of each type of Shar-Pei (meatmouth type and traditional type) were sequenced. The two traditional type Shar-Pei were sequenced at different read lengths but were aligned using the same strict criteria (allowing two mismatches per read) and therefore vary in the percentage of mapped reads as well as coverage when compared to the other individuals. Individual 7 (Table S4) was sequenced from whole genome amplified material and this may have impacted the ability to map reads and detect SNPs. This individual was not plotted in Figure 3A, but was used in downstream analyses.

Polymerase Chain Reaction (PCR) and Sanger Sequencing

All primers used were designed using Primer3 (<http://frodo.wi.mit.edu/primer3/>) [39] and are listed in Table S6. PCR and Sanger Sequencing was performed to investigate putative mutations (ten SNPs and one indel) and were carried out with 20 ng genomic DNA using AmpliTaq Gold DNA Polymerase (Applied Biosystems) following the manufacturer's instructions. The amplification of the copy number variant (CNV) breakpoints was performed with 400 ng of DNA and a Long-range PCR with Expand Long Template PCR System Mix 1 (Roche), cloned using Zero Blunt TOPO Cloning Kit (Invitrogen) and plasmid DNA prepared using QIAprep Spin Miniprep Kit (QIAGEN). PCR products and plasmids were sequenced using capillary electrophoresis 3730xl (Applied Biosystems), aligned and analyzed using CodonCode Aligner version 2.0.6 (CodonCode).

Southern blot analysis

Four micrograms of genomic DNA from each sample was digested with *Bsr*GI (New England BioLabs) and separated on a 0.7% agarose gel. A 910 bp probe (targeting CanFam 2.0 Chr13: 23,746,12–23,747,522) was used to detect the duplicated region.

Copy number assay

Estimation of copy number was performed using the comparative CT ($\Delta\Delta$ CT) relative quantification method and a calibrator animal (German Shepherd 95). The duplex reaction contained a primer limited copy number assay (CNV-E: 300 nM each of forward and reverse primers, 250 nM FAM labeled MGB probe; CNV-748: 50 nM of forward and 300 nM reverse primers, 250 nM FAM labeled MGB probe, Applied Biosystems) and a reference assay designed to C7orf28B (900 nM of forward and reverse primers, 250 nM VIC and TAMRA labeled probe, Applied Biosystems). Real Time PCR was performed in quadruplet using 10 ng of gDNA, Genotyping Master Mix (Applied Biosystems) and a 7900 HT Real Time PCR machine (Applied Biosystems). The PCR primers used and dogs evaluated can be found in Tables S4 and S1 respectively.

Fibroblast cultures

Cultures of dermal fibroblasts were established from skin samples of Shar-Pei dogs as described previously [40]. Skin samples were well shaved and cleaned with 70% EtOH/Betadine before biopsy and cell isolation. Fat tissue and blood vessels were removed from the skin and then samples were washed with PBS, cut into small fragments (0.5 cm²) and digested with dispase II solution (Boehringer Mannheim) for 16 h at 4°C. The next day, after incubation for 30 min at 37°C in the same solution, the dermis was separated from the epidermis. Washed dermal samples were chopped into 1 mm³ fragments and incubated for 140 min in 15 ml of DMEM per gram of skin containing 30 mg bacterial collagenase (Gibco), 18 mg hyaluronidase, 12 mg pronase, 1.5 mg

DNase, supplemented with bovine albumin (all from Sigma) and antibiotics. After digestion, cutaneous cells were washed with PBS and grown in a humidified atmosphere at 37°C with 5% CO₂ for two days. Medium was changed twice a week and cells were used at passages two-five.

Gene expression analysis

RNA extraction from fibroblast cultures was performed as described elsewhere [41]. 500 ng of RNA was reverse transcribed using the High-Capacity cDNA Archive Kit (Applied Biosystems) with random primers and following the manufacturer's instructions. Two assays were designed to target *HAS2* and *HAS2as* cDNA, respectively. Real Time PCR in a volume of 20 ul was performed in duplicate using SYBR Green PCR Master Mix (Applied Biosystems) and primers at 300 nM in a 7900 HT Real-Time PCR system (Applied Biosystems) with standard cycling. PCR specificity assessment was performed by adding a dissociation curve analysis at the end of the run. Each amplification run contained negative controls. Relative fold-enrichment was performed using the comparative Δ CT-method with Glucose-6-phosphate dehydrogenase (*G6PD*) for normalization.

Web resources

<http://pngu.mgh.harvard.edu/~purcell/plink/>
<http://www.codoncode.com/>
<http://genome.ucsc.edu/>
<http://maq.sourceforge.net/>
<http://frodo.wi.mit.edu/primer3/>

Supporting Information

Figure S1 Association study stratification plots prior to correction. The relationship between all individuals used in the association study was examined by plotting components C1 and C2 from the multidimensional scaling analysis. Shar-Pei individuals were grouped into two classes, cases affected by Shar-Pei Fever (FSF+A, FSF+; meatmouth type n = 24) and controls unaffected by Shar-Pei Fever (H+; traditional type n = 4 and meatmouth type n = 13). After removing two outlier cases (red circle) the genomic inflation factor was 1.2.

Found at: doi:10.1371/journal.pgen.1001332.s001 (0.47 MB TIF)

Figure S2 Comparison of the two assays for duplication detection. Two assays were developed to detect the duplications. The results from each assay for a subset of dogs tested are plotted side by side to enable comparison. The first assay, CNV-E, is targeted to the meatmouth duplication (plotted on left hand side). The second assay, CNV-748, detects both the traditional and meatmouth duplication variants (plotted right hand side). The graph shows that meatmouth type Shar-Pei also harbour the traditional duplication, however traditional type Shar-Pei (indicated with an asterisk) do not carry the meatmouth duplication. None of the control breeds sampled to date contain either duplication.

Found at: doi:10.1371/journal.pgen.1001332.s002 (1.57 MB TIF)

Figure S3 No significant relationship detected between serum hyaluronic acid (HA) concentration and copy number. Individuals have been divided based on their breed and health status. For each dog, meatmouth copy number is plotted using the left hand y-axis and serum HA levels (μ g/L) is on the right hand y-axis. All individuals with HA levels higher than 600 μ g L⁻¹ carried the duplication, although no significant correlation could be found between HA concentration and copy numbers.

Found at: doi:10.1371/journal.pgen.1001332.s003 (1.51 MB TIF)

Table S1 Individuals used in each experiment.

Found at: doi:10.1371/journal.pgen.1001332.s004 (0.07 MB PDF)

Table S2 SNPs highly associated with Familial Shar-Pei Fever on Chromosome 13. A summary of 7 SNPs on chromosome 13 that together formed a peak of association to the susceptibility of Familial Shar-Pei Fever (FSF). 41 Shar-Pei classified as affected (n = 24, FSF+ and FSF+A) and unaffected (n = 17, H+) by FSF were genotyped with an 18 K SNPs and analyzed based on 100,000 permutations. After correcting for stratification two outliers (affected by FSF) were removed and the association remained (best SNP, p = 0.01).

Found at: doi:10.1371/journal.pgen.1001332.s005 (0.06 MB PDF)

Table S3 Comparative alignment of HAS2 and HAS2 antisense.

Found at: doi:10.1371/journal.pgen.1001332.s006 (0.06 MB PDF)

Table S4 Summary statistics of the targeted resequencing. The dogs used for targeted sequencing were as follows: Control 1 = Pug, Control 2 = Napolitan Mastiff, Control 3 = Standard Poodle, Shar-Pei 1-2 are of meatmouth type and Shar-Pei 3-4 are of the traditional type. Two SNPs (Chr 13: 23,348,686 and 23,379,995) only appeared in the four sequenced Shar-Pei dogs and not in control breeds but excluded as causative for the breed-specific skin phenotype (hyaluronanosis) after further genotyping in more control breeds.

Found at: doi:10.1371/journal.pgen.1001332.s007 (0.05 MB PDF)

References

- Fraser JR, Laurent TC, Laurent UB (1997) Hyaluronan: its nature, distribution, functions and turnover. *Journal of Internal Medicine* 242: 27–33.
- Wheeler-Jones CP, Farrar CE, Pitsillides AA (2010) Targeting hyaluronan of the endothelial glycocalyx for therapeutic intervention. *Current Opinion in Investigational Drugs* 11(9): 997–1006.
- Laurent TC, Fraser JRE (1992) Hyaluronan. *FASEB J* 6: 2397–2404.
- Zanna G, Fondevila D, Bardagi M, Docampo MJ, Bassols A, et al. (2008) Cutaneous mucinosis in shar-pei dogs is due to hyaluronic acid deposition and is associated with high levels of hyaluronic acid in serum. *Vet Dermatol* 19: 314–318.
- Ramsden CA, Bankier A, Brown TJ, Cowen PSJ, Frost GI (2000) A new disorder of hyaluronan metabolism associated with generalized folding and thickening of the skin. *J of Ped* 36: 62–68.
- Weigel PH, Hascall VC, Tammi M (1997) Hyaluronan synthases. *J Biol Chem* 272: 13997–4000.
- Zanna G, Docampo MJ, Fondevila D, Bardagi M, Bassols A, et al. (2009) Hereditary cutaneous mucinosis in Shar-Pei dogs is associated with increases hyaluronan synthase-2 mRNA transcription by cultured dermal fibroblasts. *Vet Dermatol* 20: 377–382.
- Rivas AL, Tintle L, Kimball ES, Scarlett J, Quimby FW (1992) A canine febrile disorder associated with elevated interleukin-6. *Clin Immunol Immunopathol* 64: 36–45.
- Stojanov S, Kastner DL (2005) Familial autoinflammatory diseases: genetics, pathogenesis and treatment. *Curr Opin Rheumatol* 17(5): 586–599.
- Smith JM, Haigh J (1974) The hitchhiking effect of a favorable gene. *Genetic Research* 23: 23–35.
- Akey JM, Ruhe AL, Akey DT, Wong AK, Connelly CF, et al. (2010) Tracking footprints of artificial selection in the dog genome. *PNAS* 107: 1160–1165.
- Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, et al. (2007) PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 81: 559–575.
- Chao H, Spicer AP (2005) Natural Antisense mRNAs to Hyaluronan Synthase 2 Inhibit Hyaluronan Biosynthesis and Cell Proliferation. *J Biol Chem* 19: 27513–27522.
- Giuffra E, Törnsten A, Marklund S, Bongcam-Rudloff E, Chardon P, et al. (2002) A large duplication associated with dominant white color in pigs originated by homologous recombination between LINE elements flanking KIT. *Mamm Genome* 13(10): 569–77.
- Rosengren Pielberg G, Golovko A, Sundström E, Curik I, Lennartsson J (2008) A cis-acting regulatory mutation causes premature hair graying and susceptibility to melanoma in the horse. *Nat Genet* 40(8): 1104–1109.
- Salmon Hillbertz NHC, Isaksson M, Karlsson EK, Hellmén E, Pielberg GR, et al. (2007) A duplication of FGF3, FGF4, FGF9 and ORAOV1 causes the hair ridge and predisposes to dermoid sinus in Ridgeback dogs. *Nat Genet* 38(11): 1318–1320.
- Wright D, Boije H, Meadows JR, Bed'hom B, Vieaud A, et al. (2009) Copy number variation in intron 1 of SOX5 causes the pea-comb in chickens. *PLoS Genet* 5(6): 1000512. doi:10.1371/journal.pgen.1000512.

Table S5 SNPs identified in conserved elements, HAS2 and HAS2 antisense after targeted sequencing.

Found at: doi:10.1371/journal.pgen.1001332.s008 (0.10 MB PDF)

Table S6 Primers and probes used in the different experiments.

Found at: doi:10.1371/journal.pgen.1001332.s009 (0.06 MB PDF)

Acknowledgments

We thank all the dog owners, breeders, and breed clubs worldwide that have supported this study and contributed samples. We thank veterinarians and other colleagues for their help with samples including Jérôme Abadie, Laila Irene Baek, Kikka Posti, Hannes Lohi, Anne-lise Juncker, Barbara LaVere, Patricia and Harry Roach, and Jeff Vidt. We also thank Kathleen Long and Lazara Cuza for Shar-Pei photos. We thank Marie Lindersson and Kristina Larsson at the SNP Technology Platform in Uppsala (Sweden), the Broad Institute Genetic Analysis platform (US), as well as Freyja Inslund and Snaevor Sigurdsson for technical assistance with Illumina sequencing and bioinformatics.

Author Contributions

Conceived and designed the experiments: MO JRSM KT GRP FP AB ACA DLK AS ÅH EFR LA LF LT KLT. Performed the experiments: MO JRSM GRP FP JQ NT GZ MJD ACA LF. Analyzed the data: MO JRSM KT GRP FP EM JQ NT GZ EKK EBR MTW KLT. Contributed reagents/materials/analysis tools: GZ AB ACA AT ÅH LA LF LT KLT. Wrote the paper: MO JRSM LT KLT.

- Sebat J, Lakshmi B, Malhotra D, Troge J, Lese-Martin C, et al. (2007) Strong association of de novo copy number mutations with autism. *Science* 316: 445–449.
- Goldberg RL, Huff JP, Lenz ME, Glickman P, Katz R, et al. (1991) Elevated plasma levels of hyaluronan in patients with osteoarthritis and rheumatoid arthritis. *Arthritis Rheum* 34: 799–807.
- Hedin PJ, Weitoft T, Hedin H, Engström-Laurent A, Saxne T (1991) Serum concentration of hyaluronan and proteoglycans in joint disease. Lack of association. *J Rheumatol* 18: 1601–1605.
- Puré E, Assouan RK (2009) Rheostatic signaling by CD44 and hyaluronan. *Cell Signal* 21: 651–655.
- Hascall VC, Majors AK, De La Motte CA, Evanko SP, Wang A, et al. (2004) Intracellular hyaluronan: a new frontier for inflammation? *Biochim Biophys Acta* 1673: 3–12.
- Stern R, Asari AA, Sugahara KN (2006) Hyaluronan fragments: An information-rich system. *Eur J Cell Biol* 85(8): 699–715.
- Eberlein M, Schneider KA, Black KE, Collins SL, Chan-Li Y, et al. (2008) Anti-oxidant inhibition of hyaluronan fragment-induced inflammatory gene expression. *J Inflamm* 5: 20.
- Yamasaki K, Muto J, Taylor KR, Cogen AL, Audish D, et al. (2009) NLRP3/Cryopyrin is necessary for Interleukin-1 β (IL-1 β) release in response to hyaluronan, an endogenous trigger in inflammation in response to injury. *J Biol Chem* 284(19): 12762–12771.
- Kastner DL, Aksentjevich I, Goldbach-Mansky R (2010) Autoinflammatory Disease reloaded: A clinical perspective. *Cell* 140: 784–790.
- Shimada SG, Ottersen IG, Stitt JT (1994) A study of the mechanism of action of the mild analgesic dipyrone. *Agents Actions* 41: 188–192.
- deSouza GE, Cardoso RA, Melo MC, Fabricio AS, Silva VM, et al. (2002) A comparative study of the antipyretic effects of indomethacin and dipyrone in rats. *Inflamm Res* 51(1): 24–32.
- Prada J, Daza R, Chambers O, Loayza I, Huicho L (2006) Antipyretic efficacy and tolerability of oral ibuprofen, oral dipyrone and intramuscular dipyrone in children: a randomized controlled trial. *Sao Paulo Med J* 124(3): 135–140.
- DiBartola SP, Tarr MJ, Webb DM, Giger U (1990) Familial renal amyloidosis in Chinese Shar Pei dogs. *J Am Vet Assoc* 15: 483–487.
- Stridh S, Kerjaschki D, Chen Y, Rugenheimer L, Åstrand ABM, et al. (2010) Angiotensin converting enzyme inhibition blocks interstitial hyaluronan dissipation in the neonatal rat kidney via hyaluronan synthase 2 and hyaluronidase 1. *Matrix Biol*. In press.
- López A, Spracklin D, McConkey S, Hanna P (1999) Cutaneous mucinosis and mastocytosis in shar-pei. *Can Vet J* 40: 881–883.
- Miller DM (1995) The occurrence of mast cell tumors in young shar-peis. *J Vet Diagn Invest* 7: 360–363.
- Takano H, Nakazawa S, Shirata N, Tamba S, Furuta K, et al. (2009) Involvement of CD44 in mast cell proliferation during terminal differentiation. *Lab Invest* 89(4): 446–455.

35. Tanaka S (2010) Targeting CD44 in mast cell regulation. *Expert Opin Ther Targets* 14(1): 31–43.
36. Triggs-Raine B, Salo TJ, Zhang H, Wicklow BA, Natowicz MR (1999) Mutations in *HYAL1*, a member of a tandemly distributed multigene family encoding disparate hyaluronidase activities, cause a newly described lysosomal disorder, mucopolysaccharidosis IX. *Proc Natl Sci USA* 11: 6296–6300.
37. Lindblad-Toh K, Wade CM, Mikkelsen TS, Karlsson EK, Jaffe DB, et al. (2005) Genome sequence, comparative analysis and haplotype structure of the domestic dog. *Nature* 438: 803–819.
38. Heng L, Ruan J, Durbin R (2008) Mapping short DNA sequencing reads and calling variants using mapping quality scores. *Genome Res* 18: 1851–1858.
39. Rozen S, Skaletsky HO (2000) Primer3 on the WWW for general users and for biologist programmers. *Bioinformatics Methods and Protocols: Methods in Molecular Biology*. TotowaNew Jersey, USA: Humana Press. pp 365–386.
40. Serra M, Brazis P, Puigdemont A, Fondevilla D, Romano V, et al. (2007) Development and characterization of a canine skin equivalent. *Exp Dermatol* 16: 135–142.
41. Chomczynski P, Mackey K (1995) Short technical report. Modification of the TRIZOL reagent procedure for isolation of RNA from Polysaccharide-and proteoglycan-rich sources. *Biotechniques* 19(6): 942–5.